

# Sodium Hypochlorite Accident in Endodontics: An Update Review

Mohammad Sarhan AL-Zahrani\*<sup>1</sup> and Ahmad G Al-Zahrani<sup>2</sup>

<sup>1</sup>Department of Endodontic, Albaha Dental Center, King Fahad Hospital Albaha, Saudi Arabia

<sup>2</sup>Taif Dental Center Moh, Saudi Arabia

\*Corresponding author: Mohammad Sarhan AL-Zahrani, Head Department of Endodontic, Albaha Dental Center, King Fahad Hospital Albaha, Saudi Arabia, Tel: 00966543291242; E-mail: sarhan\_212@yahoo.com

Received date: 26 Nov 2015; Accepted date: 25 Dec 2015; Published date: 2 Jan 2016.

Citation: AL-Zahrani MS, AL-Zahrani AG (2016) Sodium Hypochlorite Accident in Endodontics: An Update Review. Int J Dent Oral Health 2(2): doi <http://dx.doi.org/10.16966/2378-7090.168>

Copyright: © 2016 AL-Zahrani MS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

Sodium hypochlorite accident is a very serious complication that might occur if sodium hypochlorite extrudes into the periodontal tissues. Although Sodium hypochlorite is more frequently used as endodontic irrigant. Its accident is uncommon to be encountered in the dental clinic. In this review article the sodium hypochlorite accident was investigated from different perspectives based on online review of the published papers on the same topic. Frequency, causes, clinical manifestations, complications, management and prevention were collectively reviewed in this paper.

**Keywords:** Sodium hypochlorite; Sodium hypochlorite accident; Root canal irrigant; Sodium hypochlorite complication

## Introduction

Sodium hypochlorite (NaOCl) was initially reported as an effective endodontic irrigant in 1920 [1]. Because of its antimicrobial activity, low viscosity, ability to dissolve the tissues, lubrication action and availability it is considered to be the most frequently used endodontic irrigant [2-7]. Therefore, it is used as an adjunct to mechanical debridement of the root canal system.

On the other hands, the main disadvantages of NaOCl are cytotoxicity [8]. In addition, it does not remove the smear layer without chelating agent. [6] It also alters the properties of dentin due to the dissolving action of NaOCl to the organic material such as dentin collagen [9,10]. Due to the release of chlorine, it is very toxic to vital tissues [11-13], that might lead reversible or irreversible tissue damage [14,15].

The NaOCl accident occurs when there is a considerable amount of irrigant extruded beyond the root canal space leading to tissue necrosis and induces a massive acute inflammatory response with its associated sequelae [15]. The reported cases of sodium hypochlorite accident have shown variety of associated complications, causes and different approaches for management.

The purpose of this article is to gather the scattered pieces of information about the NaOCl accident throughout the published literature, and to formulate a more decent image of sodium hypochlorite accident etiology, incidence and distribution, complication, management, and prevention.

## Action Mechanism of NaOCl

Because of the dynamic balance of sodium hypochlorite; hypochlorous acid (HOCl-), hypochlorite ions (OCl-), and hydroxyl group (OH+) will be constantly released [16].

In the presence of the above mentioned molecules different kinds of reactions occur when the sodium hypochlorite get exposed to the organic components of any cellular structure. First is the saponification reaction in which the lipids will be degraded into glycerol and lead to soap formation. Second is the neutralization reaction by which amino

acids will be neutralized by the action of NaOCl neutralizes amino acids forming water and salt. The last type of reaction is the chloramination in which amino acids will be degraded and hydrolyzed [16].

The alkalinity (pH>11) of sodium hypochlorite which is similar to the calcium hydroxide is also playing a major role in its antimicrobial and cytotoxicity activities. The high pH of NaOCl will lead to disruption of the cytoplasmic membrane and the cellular metabolism [16,17].

## Etiology

Injecting the sodium hypochlorite through the canal into the surrounding periodontal tissue is the cardinal cause of the massive tissue destruction that occurs with NaOCl accident cases. The most frequently reported mishaps that lead to NaOCl accident are root perforations, incorrect working length, and forcing the irrigation needle into narrow root canal space [15]. Other factors are also involved in the development of NaOCl accident which includes the anatomical variations, concentration of the irrigant and hypersensitivity to NaOCl.

Although the high concentration NaOCl is more efficient as antimicrobial agent, it is going to be more cytotoxic [18]. More NaOCl accidents were reported when higher concentrations of NaOCl were used. These results seem to agree with reports in the literature describing the direct relationship between the concentration and the toxicity [18,19].

It was hypothesized that sodium hypochlorite accident would occur more likely when NaOCl had direct access to a soft-tissue space, such as the buccal or infraorbital, rather than simply contacting periapical tissue [18]. In fact, this might occur more frequently when the anatomic apex of a tooth naturally fenestrated through the overlying alveolar bone or when the alveolar bone had been perforated by a disease process. The above mentioned hypothesis is supported by the increased reported cases which are associated with teeth encased in thinner cortical plate [18].

The etiology of NaOCl accident that was reported in the literature was referring to the toxic effects of sodium hypochlorite rather than to its allergic effects. Sodium hypochlorite allergic action was reported in the literature long time ago [20,21]. Another case was reported later

in which the patient developed the symptoms of allergic reaction such as hypotension and difficulty of breathing after the use of 0.5 sodium hypochlorite irrigant [22]. Allergy skin scratch test was performed which showed a positive result to NaOCl.

### **Incidence and distribution**

Sodium hypochlorite accidents were most frequently reported in the maxillary teeth (73%) compared with the mandibular teeth (21%) [18]. There was a significant difference in the region of NaOCl accident, (70%) were occurred in the molar or premolar region, whereas (30%) in the incisor or canine regions [18]. Mandibular premolar and molar teeth are encased in a denser cortical plate, and their apices are located more centrally within the body of the mandible. In contrast to this, the buccal roots of maxillary premolar and molar teeth have only a thin covering of cortical bone, which probably predisposes these teeth to NaOCl accidents [23].

Among the cases when sex was recalled, accidents were reported more frequently for females (69%) than for males (31%). This could be due to decreased bone thickness and density of the female patient [18].

### **Clinical manifestation**

When NaOCl accident occurs the patient usually developed a rapid intra and extra oral swelling which might be edematous, hemorrhagic or both due to the acute inflammatory process [12]. The swelling might extends beyond the site of the affected tooth [13,14]. The most pathognomonic sign is the sudden pain which will be felt during irrigation of the root canal space [6,15]. Ecchymosis and bruising might also develop intra-orally and/or extra-orally due to bleeding into the interstitial tissues [8,12]. Moreover, the adjacent mucosa might get necrotized due to the chemical burn of NaOCl [24]. Tissue necrosis may appear within minutes or might occur hours or days later [8,25].

The upper airway might be affected in some cases. This could happen if the submental and sublingual spaces will be involved [26]. The signs of upper air way obstruction include stridor, elevation of the floor of the mouth, labored breathing and declining oxygen saturation [26].

If the reaction was due to allergic reaction to NaOCl the clinical manifestation will includes the usual symptoms of allergy. These might include hypotension, shortness of breath, edema, wheezing and urticaria [22].

### **Complications**

#### **Scar formation**

When tissue necrosis occurred due to the chemical burn of NaOCl, secondary intention healing process might happen that will lead to scar formation [27].

#### **Sinusitis**

Involvement of the maxillary sinus during endodontic therapy most probably leads to acute sinusitis. Hence, congestion of the maxillary sinus is cardinal sign if the sinus was affected by the irrigant [28].

In the reported cases of affected sinuses with NaOCl the symptoms are ranging from nothing more than smell and taste of sodium hypochlorite to heavily congested sinuses and severe pain [28,29].

#### **Neurological complications**

Parasthesia of the affected tissues might also develop due to involvement of the sensory nerves by the chemical burn of the sodium hypochlorite [30].

If the motor nerves were affected by NaOCl accident then the motor function of the supplied muscles will be affected also [31].

Fortunately, the affected nerves usually will restore their normal function, but in different period of time that might extends to several months [30].

### **Upper airway obstruction**

Life threatening air way obstruction has been reported as sequelae of NaOCl extrusion. In the reported case of the upper air way obstruction due to NaOCl accident the cellulitis and swelling spread rapidly to involve the submental and sublingual spaces and resulting in elevation of the floor of the mouth. Then the patient started to exhibit signs of upper airway obstruction [26].

Upper air way obstruction could results also if the solution was inhaled or ingested [30,14].

### **Management**

Unfortunately, low level evidences are available for the management of sodium hypochlorite accident complications such as expert opinion or case reports. Thus, there is no standard treatment protocol has been documented; this could be because these complications are rare and uncommon.

The first step for management is to explain the case to the patient. This step plays a crucial role in dealing with the consequences of this kind of accident. Also, documents all the details of the incident including irrigation needle type, the use of rubber dam, working length, volume and concentration of NaOCl. Take clinical pictures to support these notes.

Usually conservative and palliative management of NaOCl accident is advocated [32]. However, treatment will be determined by the severity of the case [33].

Pain could be managed using long acting local anesthetic and analgesic such as non steroidal anti-inflammatory drugs and paracetamol. Flexible prescription by alternating ibuprofen and paracetamol at four hours interval might be effective for severe pain management [1].

It is recommended to prescribe a prophylactic antibiotic to prevent the secondary infection that might develop due to tissue necrosis or hematoma in the affected tissues [31].

The use of steroids might be also recommended to control the process of acute inflammation [33,34]. Several clinical studies reported that administration of dexamethasone is effective in minimizing postoperative pain and swelling after endodontic therapy or flare-up cases [35-37].

Advise the patient to use cold compression for the first day to control the swelling. Surgical drainage might be also needed to relief the congested tissue [31].

In case of upper airway obstruction urgent intervention and hospitalization is recommended. Intubation and administration of intravenous steroids and antihistamines will be necessary [26].

In the cases of maxillary sinus involvement it might be necessary to drain the sinus surgically [28]. However if the sinus doesn't become congested, irrigation of the sinus through the root canal using distilled water or saline might be enough [29].

### **Prevention**

As mentioned before, these complications are rare. Nonetheless, the risk of accident could be minimized by implementation of the preventive measures when performing endodontic therapy.

Control of Substances Hazardous to Health Regulations (COSHH) is recommending that the exposure of the patient to NaOCl must be kept as low as reasonable practicable. So the use of rubber dam is a must to isolate the tooth for root canal treatment [21].

It is very important that clinician must investigate thoroughly the presence of any predisposing risk factor that might lead to development of NaOCl accident such as perforations, resorption, immature apices or any other conditions [29].

The side vented needle is advocated for root canal. The end vented needles carry risks of irrigant extrusion [15,38]. The needle must not binds to the walls of the canal, but to be kept loose while irrigating. Moreover, it should be kept in at least two mm distance from the apical terminus [15]. The irrigant should be introduced with low pressure to prevent forcing it beyond the apex [15]. Even though, the use of negative pressure irrigation system such as EndoVac system to the full (WL) was principally proven to minimize the risk of extrusion of root canal irrigant [39].

Care should be taken in the management of teeth with immature root and open apices to prevent the irrigant from seepage beyond the root apex. It has been suggested to limit the use of NaOCl irrigation to the coronal 2/3 of the root with open apex [18].

It is very important to be aware of the possibility of sodium hypochlorite allergic reactions and take measures to prevent occurrence.

#### The questions usually asked in a medical history report are:

Are you sensitive to household bleach?

Did you experience any sensitivity when swimming in a chlorinated pool?

Are you allergic to Intravenous Pyelogram dye? (Dye contains iodine).

Some authors said that this allergy may indicate that the patient is more likely to be allergic to other substances such as chlorine, [40] and eventually will be susceptible for NaOCl accident. Allergy skin scratch test is recommended particularly in cases with expected NaOCl allergy before endodontic treatment [38].

#### Acknowledgement: nil

#### References

- Crane A B (1920) A practicable root canal technique. Philadelphia: Lea and Febinger.
- European Society of Endodontology (2006) Quality guidelines for endodontic treatment: consensus report of the European Society of Endodontology. *Int Endod J* 39: 921-30.
- Grandini S, Balleri P, Ferrari M (2002) Evaluation of Glyde File Prep in combination with sodium hypochlorite as a root canal irrigant. *J Endod* 28: 300-3.
- Averbach RE, Kleier DJ (2006) Clinical update on root canal disinfection. *Compend Contin Educ Dent* 27: 284-9.
- Regan JD, Fleury AA (2006) Irrigants in non-surgical endodontic treatment. *J Ir Dent Assoc*. 52: 84-92.
- McComb D, Smith DC (1975) A preliminary scanning electron microscopic study of root canals after endodontic procedures. *J Endod* 1: 238-42.
- Ercan E, Ozekinci T, Atakul F, Gül K (2004) Antibacterial activity of 2% chlorhexidine gluconate and 5.25% sodium hypochlorite in infected root canal: in vivo study. *J Endod* 30: 84-7.
- Gomes BP, Ferraz CC, Vianna ME, Berber VB, Teixeira FB, et al. (2001) In vitro antimicrobial activity of several concentrations of sodium hypochlorite and chlorhexidine gluconate in the elimination of *Enterococcus faecalis*. *Int Endod J* 34: 424-8.
- Sim TP, Knowles JC, Ng YL, Shelton J, Gulabivala K (2001) Effect of sodium hypochlorite on mechanical properties of dentine and tooth surface strain. *Int Endod J* 34: 120-32.
- Grigoratos D, Knowles J, Ng YL, Gulabivala K (2001) Effect of exposing dentine to sodium hypochlorite and calcium hydroxide on its flexural strength and elastic modulus. *Int Endod J* 34: 113-9.
- Zehnder M, Kosicki D, Luder H, Sener B, Waltimo T (2002) Tissue-dissolving capacity and antibacterial effect of buffered and unbuffered hypochlorite solutions. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 94: 756-62.
- Lamers AC, van Mullem PJ, Simon M (1980) Tissue reactions to sodium hypochlorite and iodine potassium iodide under clinical conditions in monkey teeth. *J Endod* 6: 788-92.
- Gernhardt CR, Eppendorf K, Kozłowski A, Brandt M (2004) Toxicity of concentrated sodium hypochlorite used as an endodontic irrigant. *Int Endod J* 37: 272-80.
- Witton R, Henthorn K, Ethunandan M, Harmer S, Brennan PA (2005) Neurological complications following extrusion of sodium hypochlorite solution during root canal treatment. *Int Endod J* 38: 843-8.
- Hülsmann M, Hahn W (2000) Complications during root canal irrigation-literature review and case reports. *Int Endod J* 33: 186-93.
- Estrela C, Estrela CR, Barbin EL, Spanó JCE, Marchesan MA, et al. (2002) Mechanism of action of sodium hypochlorite. *Braz Dent J* 13: 113-117.
- Estrela C, Sydney GB, Bammann LL, Júnior FO (1995) Mechanism of action of calcium and hydroxyl ions of calcium hydroxide on tissue and bacteria. *Braz Dent J* 6: 85-90.
- Kleier DJ, Averbach RE, Mehdipour O (2008) The Sodium Hypochlorite Accident: Experience of Diplomates of the American Board of Endodontics. *J Endo* 34: 1346-1350.
- Pashley EL, Bridesong NL, Bowman K, Pashley DH (1985) Cytotoxic effects of sodium hypochlorite on vital tissue. *J Endod* 11: 525-8.
- Sulzberger M B (1940) Dermatologic allergy: an introduction in the form of a series of lectures. Springfield, IL, USA: Charles C. Thomas.
- Cohen S, Burns R (1984) Pathways of the pulp. 3rd edn. pp 441-442. St Louis, MO, USA: CV Mosby.
- Caliskan M K, Turkun M, Alper S (1994) Allergy to sodium hypochlorite during root canal therapy: a case report. *Int Endod J* 27: 163-167.
- Vertucci FJ, Haddix JE, Britto LR (2006) Tooth morphology and access preparation. In: Cohen S, Hargreaves KM, eds. Pathways of the Pulp. 9th ed. St. Louis: Mosby Inc, 197-9.
- Sjogren U, Figdor D, Persson S, Sundqvist G (1997) Influence of infection at the time of root filling on the outcome of endodontic treatment of teeth with apical periodontitis. *Int Endod J* 30: 297-306.
- Shuping GB, Ørstavik D, Sigurdsson A, Trope M (2000) Reduction of intracanal bacteria using nickel-titanium rotary instrumentation and various medications. *J Endod* 26: 751-5.
- Al-Sebaei MO, Halabi OA, El-Hakim EI (2015) sodium hypochlorite accident resulting in life-threatening airway obstruction during root canal treatment: a casre report, *Clinical, Cosmetic and Investigational Dentistry* 7(41-44).
- Mehra P, Clancy C, Wu J (2000) Formation of a facial hematoma during endodontic therapy. *J Am Dent Assoc* 131: 67-71.
- Kavanagh C P, Taylor J (1998) Inadvertent injection of sodium hypochlorite into the maxillary sinus. *Br Dent J* 185: 336-337.
- Ehrich DG, Brian JD Jr, Walker WA (1993) Sodium hypochlorite accident: inadvertent injection into the maxillary sinus. *J Endod* 19: 180-182.
- Becking AG (1991) Complications in the use of sodium hypochlorite during endodontic treatment. *Oral Surg Oral Med Oral Path* 71: 346-348.

31. Ziegler DS, Bent GP (2001) Upper airway obstruction induced by a caustic substance found responsive to nebulised adrenaline. *J Paediatr Child Health* 37: 524-525.
32. Hales JJ, Jackson CR, Everett AP, Moore SH (2001) Treatment protocol for the management of a sodium hypochlorite accident during endodontic therapy. *Gen Dent* 49: 278-81.
33. Gatot A, Arbelle J, Leiberman A, Yanai-Inbar I (1991) Effects of sodium hypochlorite on soft tissues after its inadvertent injection beyond the root apex. *J Endodon* 17: 573-4.
34. Czerwinski AW, Czerwinski AB, Whitsett TL, Clark ML (1972) Effects of a single large intravenous injection of dexamethasone. *Clin Pharmacol Ther* 13: 638-642.
35. Liesinger A, Marshall FJ, Marshall JG (1993) Effect of variable doses of dexamethasone on post treatment endodontic pain. *J Endod* 19: 35-39.
36. Stewart G (1962) Combined use of an antibiotic and a corticosteroid for postoperative sequelae in endodontic practice. *J Dent Med* 17: 142-146.
37. Tsesis I, Fuss Z, Lin S, Tilinger G, Peled M (2003) Analysis of postoperative symptoms following surgical endodontic treatment. *Quintessence Int* 34: 756-60.
38. Kaufman AY, Keila S (1989) Hypersensitivity to sodium hypochlorite. *J Endod* 15: 224-226.
39. Mitchell RP, Baumgartner JC, Sedgley CM (2011) Apical extrusion of sodium hypochlorite using different root canal irrigation systems. *J Endod* 37: 1677-1681.
40. Shehadi WH (1975) Adverse reactions to intravascularly administered contrast media. A comprehensive study based on a prospective survey. *Am J Roentgenol Radium Ther Nucl Med* 124: 145-52.