

Challenges to Treat Interferon Resistance in Hepatitis C Virus Infected Patients

Haroon Hamed M¹, Shafiqur Rehman² and Ishtiaq Qadri^{3*}

¹Department of Biological Sciences, Faculty of Science King Abdul-Aziz University, Jeddah

²Department of Microbiology and Molecular Genetics, University of the Punjab, New Campus, Lahore Pakistan

³King Fahd Medical Research Center, King Abdul Aziz University, Jeddah 21589 Saudi Arabia

Corresponding author: Ishtiaq Qadri, King Fahd Medical Research Center, King Abdul Aziz University, Jeddah 21589, Saudi Arabia, Tel: +96-65-35168434; Fax: +96-62-6952076; E-mail: ishtiaqmq@yahoo.com, ishtiaq80262@yahoo.com

Received date: 27 May 2016; **Accepted date:** 03 Jun 2016; **Published date:** 09 Jun 2016.

Citation: Hamed HM, Rehman S, Qadri I (2016) Challenges to Treat Interferon Resistance in Hepatitis C Virus Infected Patients. *J Mol Med Clin Appl* 1(1): doi <http://dx.doi.org/10.16966/2575-0305.e102>

Copyright: © 2016 Hamed HM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Keywords: Hepatitis C virus (HCV); Interferon (IFN); Protein kinase R (PKR); Interferon stimulatory genes (ISGs); Suppressor of cell signaling (SOCS)

The host response to virus is initiated by interferon α/β (IFN) production which is the baseline of immune resistance against viral pathogens. Hepatitis C virus (HCV) infects about 170 million persons globally [1]. The major obstacle in the treatment of HCV is interferon resistance associated with each genotype of HCV, particularly the HCV GT1 and GT4. HCV eludes the host reaction over a multifaceted permutation of virus-host relations that interrupt intracellular signaling pathways and diminish the antiviral actions of IFN. Innate and adaptive immunity related to host responses and its regulation results the spread and replication of HCV. Consequently, a major challenge to the treatment responses is genetic diversity of virus and its linkage with initiation of chronic liver disease. Studies have revealed several host traits (gender, age, ethnicity, insulin resistance, obesity, alcohol intake, HIV-1 infection, degree of liver fibrosis and cirrhosis) with IFN responsiveness [2,3]. Intensive research efforts are desired to elaborate the molecular mechanisms involved in the IFN non-responsiveness and of ascertaining its predictors that could be used to plan a suitable cure regimen. Factors that may contribute to the interferon resistance are multi-factorial but clearly 1) virus-host interaction, 2) IFN signaling cascade, 3) virus mutations, 4) immune responses, 5) host's genetic makeup may contribute for the resistance phenotype.

Viral and Host Interaction

HCV infected chimpanzees have revealed a sturdy orientation of Interferon stimulating genes (ISGs) during the early days of the infection [4]. The exact mechanisms of IFN sensitivity are not entirely understood but possibly comprise the IFN inhibition via theroetic acid inducible gene I(RIG-I) pathway via the cleavage of MAVS (mitochondrial antiviral signalling protein) by the NS3 protease [5]. IL28 [6] and KIR2DS [7] polymorphisms play a vital role in the retort of chronic HCV to therapy.

Role of pegIFN α , ribavirin and IFN λ signaling

IFN α -based therapies showed resistance in the patients with stimulated endogenous IFN system [8]. The most precise extrapolation of the pegIFN α and ribavirin reaction is accomplished by the expression of ISGs of the liver biopsies [9]. IFN λ could surge high virological response in individuals with a pre-activated endogenous IFN response as USP18 (Ubl carboxyl-terminal hydrolase) does not inhibit the IFN λ signaling [10].

Viral genetic heterogeneity

The viral regions of HCV can contribute to IFN resistance via the articulation of various HCV regions like core, E1, E2, NS5A, and NS5B

[11]. The SOCS-3 expression might be induced by core that quashes JAK-STAT signalling pathways [12]. Manifestation of the PIAS (protein inhibitor of activated STAT) is tempted by HCV proteins which is probably arbitrated by PP2A (protein phosphatase 2A) signaling and STAT demethylation [13] resulting the STAT1 blockage. High level of IL-8 exhibited in HCV infected individuals [14]. Mutations at amino acids 70 and 91 of core deliberateresistance to IFN- α , related with a decline in IFN-tempted phosphorylation of STAT1 and STAT2 and expression of ISGs [15]. Some regions linked with IFN- α and ribavirin sensitivity have been documented within core region known as ISDR (a.a. 2209- 2248) and IRRDR (a.a. 2334-2379) [16].

IFN- α resistance was observed in recombinant HCV genotypes of 1a and 3a identified substitutions of amino acids at position 414 of E2 and positions 345 & 348 of E1 [17]. The interface between E2 and PKR could be one contrivance by which HCV avoids the interferon [18]. Serine phosphorylation in E2 gene of HCV GT1 had expressed a role in interferon resistance [19]. Mutations related to treatment response have been reported in NS5A region which include ISDR, residues 2209 to 2248, and IRRDR, residues 2334 to 2379, in HCV genotype 1 [20,21]. These mutations were mainly observed in IFN resistant patients. Both core and NS5A express changes during the treatment so impact the therapy consequences [22]. *In vitro* study of HCV replicon cell lines, substitutions in NS3, NS4B, NS5A, and NS5B were linked with IFN non responsiveness [23]. In another study, overexpression of NS5A GT1 showed least IFN response as compared to GT-3 over sturdier binding to STAT1 [24].

Host Immune Responses associated with Interferon Resistance

The development of persistent HCV infection is the result of virus wining the fight from host immune response. During the persistent infection, interferon being a component of innate immune response might have activated at their level best to activate the effective adaptive immune response to infection clearance. However, due to a plethora of reasons (viral mutations and host genetic variation) leads to activation of non-HCV clearance immune response even with the presence of high interferon concentration [8].

Future Directions

The critical issues related to the IFN resistance need to be elucidated in more detail. The role of endogenous IFN λ s, and molecular association between ISGs and *IFNL3* needs to be defined for further understandingthe

mechanism of HCV interference with IFN signalling and different type of ISG [2]. The hepatologists and virologists need to be vigilant to define the emerging extend of directly-acting antiviral (DAAs) resistance during treatment. This combination of effective next generation of DAAs with new class of IFN λ s should produce good results. The IFN resistance via weak immune responses will help in the design of therapy with chemokines and cytokines co-stimulation of ISG pathways in selected patients.

References

1. Shepard CW, Finelli L, Alter MJ (2005) Global epidemiology of hepatitis C virus infection. *Lancet Infect Dis* 5: 558-567.
2. Heim MH (2013) 25 years of interferon-based treatment of chronic hepatitis C: an epoch coming to an end. *Nat Rev Immunol* 13: 535-542.
3. Qashqari H, Al-Mars A, Chaudhary A, Abuzenadah A, Damanhouri G, et al. (2013) Understanding the molecular mechanism(s) of hepatitis C virus (HCV) induced interferon resistance. *Infect Genet Evol* 19: 113-119.
4. Major ME, Dahari H, Mihalik K, Puig M, Rice CM, et al. (2004) Hepatitis C virus kinetics and host responses associated with disease and outcome of infection in chimpanzees. *Hepatology* 39: 1709-1720.
5. Meylan E, Curran J, Hofmann K, Moradpour D, Binder M, et al. (2005) Cardif is an adaptor protein in the RIG-I antiviral pathway and is targeted by hepatitis C virus. *Nature* 437: 1167-1172.
6. López-Rodríguez R, Hernández-Bartolomé Á, Borque MJ, Rodríguez-Muñoz Y, Martín-Vílchez S, et al. (2013) Polymorphisms in histone deacetylases improve the predictive value of IL-28B for chronic hepatitis C therapy. *Genes Immun* 14: 317-324.
7. Keane C, O'Shea D, Reiberger T, Peck-Radosavljevic M, Farrell G, et al. (2013) Variation in both IL28B and KIR2DS3 genes influence pegylated interferon and ribavirin hepatitis C treatment outcome in HIV-1 co-infection. *PLoS One* 8: e66831.
8. Sarasin-Filipowicz M, Oakeley EJ, Duong FH, Christen V, Terracciano L, et al. (2008) Interferon signaling and treatment outcome in chronic hepatitis C. *Proc Natl Acad Sci U S A* 105: 7034-7039.
9. Dill MT, Duong FH, Vogt JE, Bibert S, Bochud PY, et al. (2011) Interferon-induced gene expression is a stronger predictor of treatment response than IL28B genotype in patients with hepatitis C. *Gastroenterology* 140: 1021-1031.
10. Makowska Z, Duong FH, Trincucci G, Tough DF, Heim MH, et al. (2011) Interferon-beta and interferon-lambda signaling is not affected by interferon-induced refractoriness to interferon-alpha *in vivo*. *Hepatology* 53: 1154-1163.
11. Xu J, Liu S, Xu Y, Tien P, Gao G, et al. (2009) Identification of the nonstructural protein 4B of hepatitis C virus as a factor that inhibits the antiviral activity of interferon-alpha. *Virus Res* 141: 55-62.
12. Alexander WS (2002) Suppressors of cytokine signalling (SOCS) in the immune system. *Nat Rev Immunol* 2: 410-416.
13. Duong FH, Filipowicz M, Tripodi M, La Monica N, Heim MH, et al. (2004) Hepatitis C virus inhibits interferon signaling through up-regulation of protein phosphatase 2A. *Gastroenterology* 126: 263-277.
14. Polyak SJ, Khabar KS, Rezeiq M, Gretch DR (2001) Elevated levels of interleukin-8 in serum are associated with hepatitis C virus infection and resistance to interferon therapy. *J Virol* 75: 6209-6211.
15. El-Shamy A, Kim SR, Ide YH, Sasase N, Imoto S, et al. (2012) Polymorphisms of hepatitis C virus non-structural protein 5A and core protein and clinical outcome of pegylated-interferon/ribavirin combination therapy. *Intervirology* 55: 1-11.
16. Fukuhara T, Taketomi A, Okano S, Ikegami T, Soejima Y, et al. (2010) Mutations in hepatitis C virus genotype 1b and the sensitivity of interferon-ribavirin therapy after liver transplantation. *J Hepatol* 52: 672-680.
17. Serre SB, Krarup HB, Bukh J, Gottwein JM (2013) Identification of alpha interferon-induced envelope mutations of hepatitis C virus *in vitro* associated with increased viral fitness and interferon resistance. *J Virol* 87: 12776-12793.
18. Chayama K, Suzuki F, Tsubota A, Kobayashi M, Arase Y, et al. (2000) Association of amino acid sequence in the PKR-eIF2 phosphorylation homology domain and response to interferon therapy. *Hepatology* 32: 1138-1144.
19. Afzal S, Idrees M, Ali M, Ilyas M, Hussain A, et al. (2011) Envelope 2 protein phosphorylation sites S75 & 277 of hepatitis C virus genotype 1a and interferon resistance: a sequence alignment approach. *Virology* 41: 71.
20. El-Shamy A, Shoji I, El-Akel W, Bilasy SE, Deng L, et al. (2012) NS5A sequence heterogeneity of hepatitis C virus genotype 4a predicts clinical outcome of pegylated-interferon-ribavirin therapy in Egyptian patients. *J Clin Microbiol* 50: 3886-3892.
21. El-Shamy A, Shoji I, Kim SR, Ide Y, Imoto S, et al. (2012) Sequence heterogeneity in NS5A of hepatitis C virus genotypes 2a and 2b and clinical outcome of pegylated-interferon/ribavirin therapy. *PLoS One* 7: e30513.
22. Kozuka R, Enomoto M, Hai H, Ogawa T, Nakaya M, et al. (2012) Changes in sequences of core region, interferon sensitivity-determining region and interferon and ribavirin resistance-determining region of hepatitis C virus genotype 1 during interferon-alpha and ribavirin therapy, and efficacy of retreatment. *Hepatol Res* 42: 1157-1167.
23. Namba K, Naka K, Dansako H, Nozaki A, Ikeda M, et al. (2004) Establishment of hepatitis C virus replicon cell lines possessing interferon-resistant phenotype. *Biochem Biophys Res Commun* 323: 299-309.
24. Kumthip K, Chusri P, Jilg N, Zhao L, Fusco DN, et al. (2012) Hepatitis C virus NS5A disrupts STAT1 phosphorylation and suppresses type I interferon signaling. *J Virol* 86: 8581-8591.