

Advances in Nanomedicine for Treatment of Stroke

Medha D Joshi*

Chicago College of Pharmacy, Department of Pharmaceutical Sciences, Midwestern University, Downers Grove, IL, 60515, USA

Corresponding author: Dr. Medha D Joshi, Department of Pharmaceutical Sciences, Midwestern University, Chicago College of Pharmacy, Alumni Hall 365, 555 31st Street, Downers Grove, IL 60515, USA, Tel: +1-630-515-6963 (office); Fax: +1-630-515-6958; **E-mail:** mjoshi@midwestern.edu

Stroke continues to be the most serious and debilitating neurological disorder in the United States. With a new event every 40 seconds and a victim every 4 minutes, it is the fifth leading cause of death and number one cause of disability. A recent update from the Stroke Statistics Subcommittee of the American Heart Association reported that about 795,000 people in the US experience a new or recurrent stroke each year [1]. Ischemic stroke (blockage of a blood vessel supplying the brain) contributes 80% of the total, and hemorrhagic stroke (bleeding into or around the brain) makes up the rest. National Institutes of Health (NIH) estimate that stroke costs exceed \$73 billion in US healthcare dollars per year [2]. Despite the severity of this condition, the only available FDA approved pharmacological treatments for ischemic stroke are: endovascular recanalization therapy and intravenous administration of recombinant tissue plasminogen activator (t-PA), which dissolves the clot and restores blood flow to the brain [3]. The limitation of the endovascular recanalization procedure is 1) only limited population with large vessel occlusions and salvageable brain tissue actually benefits from the therapy; 2) requirement of significant resources in both equipment and trained personnel to perform the surgery [4]. The treatment with t-PA is complicated by: 1) a relatively short "window of opportunity" of only 3-4.5 hours between infarct and treatment; 2) an increased risk for subarachnoid hemorrhage; and 3) neurotoxicity. As a result, only a few patients receive (3-8.5%) and benefit (1-2%) from t-PA treatment. Neither technique addresses cellular damage to brain tissue adjacent to the infarcted area, known as the stroke penumbra. More than 1,000 molecules broadly classified as neuroprotective, aiming to stop or slow the secondary damage associated with the ischemic cascade following stroke, have shown promise in the initial stages of research but have failed to demonstrate efficacy in clinical studies because of inadequate efficacy or side effects [5]. A new approach is therefore needed urgently, one which has the potential to address both the restoration of blood flow and attenuation of secondary damage to the penumbral area [5].

Recently, there has been considerable interest in the application of nanomedicine to the treatment of stroke [6-12]. The prime reasons behind their popularity are 1) feasibility of the nanomedicine to decrease neurotoxicity by decreasing the dose; 2) ability to cross the blood brain barrier and ability to accumulate in the ischemic region [13]; 3) target the clot by means of ligand based targeting; 4) release the cargo under specific stimuli [14]. Liposomes have been at the forefront of these investigations followed by polymeric nanoparticles owing to their approval for clinical use, biodegradability and biocompatibility. Other nanocarriers such as platinum [15], ceria [16] and carbon nanotubes [17] have been also investigated for their free radical scavenging properties. Various pharmaceutical actives have been investigated including: thrombolytics [18] such as t-PA [19], streptokinase [20,21] and plasmin

Received date: 06 May 2016; **Accepted date:** 07 May 2016; **Published date:** 12 May 2016.

Citation: Joshi MD (2016) Advances in Nanomedicine for Treatment of Stroke. *Int J Nanomed Nanosurg* 2(3): doi <http://dx.doi.org/10.16966/2470-3206.113>

Copyright: © 2016 Joshi MD. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

[22]; hemoglobin [23-28]; tacrolimus (FK506) [13,29-32]; superoxide dismutase [33]; asialo-erythropoietin [31,34]; IRL-1620 [35]; fasudil [36]; bioactive gases such as nitric oxide [37], Xenon [38]; Nerve Growth Factor (NGF) [39]; citicoline [40,41]; ZL006 [42,43]; antioxidant enzymes [44-46]; luteolin [47]; angiogenic peptides [48] and siRNA [49]. In an animal model of permanent middle cerebral artery occlusion (pMCAO), intravenously-administered 100 nm liposomes [13,29] were found to accumulate in the ischemic core and penumbra region when administered 1-2 hr after occlusion despite reduction in the cerebral blood perfusion as confirmed by fluorescence imaging and positron emission tomography (PET). Liposomes have also been surface modified with polyethylene glycol (PEG) coating to improve their circulation time and to improve their target specificity, by adding moieties such as transferin receptor [41,43,50,51], Fas ligand antibody [52].

A group of researchers are studying the delivery of medical gases using echogenic liposomes [19,22,37,38,53-56]. Echogenic liposomes are liposomes that encapsulate both gas and fluid in their phospholipid bilayer and act as contrast agent for ultrasound. Thrombolytics such as t-PA [55] and plasmin [22] have been encapsulated in echogenic liposomes allowing local delivery with minimal systemic exposure. Combination approaches [57,58] such as delivering antioxidant/anti-inflammatory agents with t-PA have been reported to significantly improve the treatment outcomes. Interestingly, however, when an anti-oxidant was encapsulated into and delivered by three different nanocarriers - PEGylated liposomes or polymeric nanoparticles made up of PLA and PLGA with or without targeting moiety - no significant difference was observed in infarct volume reduction [44], indicating the probability that amount of drug encapsulated in the nanocarrier is of greater importance than the choice of nanocarrier or the targeting moiety. Another relatively new area that is gaining interest is in the arena of nano-theranostics [59-61] where diagnosis and therapy is carried out simultaneously using the same nanomedicines. Despite these excellent advances in nanomedicine research that has led to increase in safety and efficacy of neuroprotective and neuroreparative agents that are used in the treatment of stroke, a clinically functioning product is still lacking.

Nanomedicine has great potential in the treatment of stroke. It has been established that nanomedicines can deliver drugs across brain for efficacious reduction in infarct volume and reduce the toxicity of treatment with t-PA when compared to conventional delivery systems. Further work, however, is needed in order to translate these advances into improved clinical outcomes. Effective nanocarrier design to improve drug encapsulation and release, rather than targeting, following STAIR guidelines [62] to evaluate efficacy of the nanomedicine in animal models is needed to improve the chances of nanomedicine being developed into a clinically useful treatment for stroke.

Acknowledgments

Author would like to acknowledge the assistance provided by E. Langan, Department of Pharmaceutical Sciences, and Midwestern University, IL, USA in preparing this opinion article.

References

- Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, et al. (2016) Heart Disease and Stroke Statistics-2016 Update: A Report From The American Heart Association. *Circulation* 133: e38-e360.
- Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, et al. (2010) Heart Disease and Stroke Statistics-2010 Update. *Circulation* 121: e46-e215.
- Meyers PM, Schumacher HC, Connolly ES Jr, Heyer EJ, Gray WA, et al. (2011) Current Status of Endovascular Stroke Treatment. *Circulation* 123: 2591-2601.
- Appireddy RMR, Demchuk AM, Goyal M, Menon BK, Eesa M, et al. (2015) Endovascular Therapy for Ischemic Stroke. *J Clin Neurol* 11: 1-8.
- Ginsberg MD (2009) Current status of neuroprotection for cerebral ischemia: synoptic overview. *Stroke* 40: S111-S114.
- Kubinova S, Sykova E (2010) Nanotechnology for treatment of stroke and spinal cord injury. *Nanomedicine* 5: 99-108.
- Lanza GM, Marsh JN, Hu G, Scott MJ, Schmieder AH, et al. (2010) Rationale for a Nanomedicine Approach to Thrombolytic Therapy. *Stroke* 41: S42-S44.
- Bhaskar S, Tian F, Stoeger T, Kreyling W, de la Fuente JM, et al. (2010) Multifunctional Nanocarriers for diagnostics, drug delivery and targeted treatment across blood-brain barrier: Perspectives on tracking and neuroimaging. *Part Fibre Toxicol* 7: 3.
- Kim KS, Khang G, Lee D (2011) Application of nanomedicine in cardiovascular diseases and stroke. *Curr Pharm Des* 17: 1825-1833.
- Nair SB, Dileep A, Rajanikant GK (2012) Nanotechnology based diagnostic and therapeutic strategies for neuroscience with special emphasis on ischemic stroke. *Curr Med Chem* 19: 744-756.
- Sharma HS (2011) Blood-CNS barrier, neurodegeneration and neuroprotection: Recent therapeutic advancements and nano-drug delivery. *J Neural Transm (Vienna)* 118: 3-6.
- Gaudin A, Andrieux K, Couvreur P (2015) Nanomedicines and stroke: Toward translational research. *J Drug Deliv Sci Technol* 30: 278-299.
- Ishii T, Fukuta T, Agato Y, Oyama D, Yasuda N, et al. (2013) Nanoparticles accumulate in ischemic core and penumbra region even when cerebral perfusion is reduced. *Biochem Biophys Res Commun* 430: 1201-1205.
- Fukuta T, Asai T, Oku N (2015) Development of a liposomal drug delivery system for the treatment of ischemic stroke. *Drug Deliv Sys* 30: 309-316.
- Takamiya M, Miyamoto Y, Yamashita T, Deguchi K, Ohta Y, et al. (2012) Strong neuroprotection with a novel platinum nanoparticle against ischemic stroke- and tissue plasminogen activator-related brain damages in mice. *Neuroscience* 221: 47-55.
- Kim CK, Kim T, Choi IY, Soh M, Kim D, et al. (2012) Ceria nanoparticles that can protect against ischemic stroke. *Angew Chem Int Ed Engl* 51: 11039-11043.
- Lee HJ, Park J, Yoon OJ, Kim HW, Lee do Y, et al. (2011) Amine-modified single-walled carbon nanotubes protect neurons from injury in a rat stroke model. *Nat Nanotechnol* 6: 121-125.
- Elbayoumi TA, Torchilin VP (2008) Liposomes for targeted delivery of antithrombotic drugs. *Expert Opin Drug Deliv* 5: 1185-1198.
- Laing ST, Moody MR, Kim H, Smulevitz B, Huang SL, et al. (2012) Thrombolytic efficacy of tissue plasminogen activator-loaded echogenic liposomes in a rabbit thrombus model. *Thromb Res* 130: 629-635.
- Leach JK, O'Rear EA, Miao Y, Johnson AE, Patterson ES (1999) Thrombolysis in a rabbit stroke model using liposomal-encapsulated streptokinase. In: *Proceedings of Annual International Conference of the IEEE Engineering in Medicine and Biology* 776.
- Leach JK, Patterson E, O'Rear EA (2004) Improving thrombolysis with encapsulated plasminogen activators and clinical relevance to myocardial infarction and stroke. *Clin Hemorheol Microcirc* 30: 225-228.
- Kandadai MA, Meunier JM, Hart K, Holland CK, Shaw GJ (2014) Plasmin-Loaded Echogenic Liposomes for Ultrasound-Mediated Thrombolysis. *Transl Stroke Res* 6: 78-87.
- Fukumoto D, Kawaguchi AT, Haida M, Yamano M, Ogata Y, et al. (2009) Liposome-encapsulated hemoglobin reduces the size of cerebral infarction in rats: Effect of oxygen affinity. *Artif Organs* 33: 159-163.
- Kakehata J, Yamaguchi T, Togashi H, Sakuma I, Otani H, et al. (2010) Therapeutic potentials of an artificial oxygen-carrier, liposome-encapsulated hemoglobin, for ischemia/reperfusion-induced cerebral dysfunction in rats. *J Pharmacol Sci* 114: 189-197.
- Kawaguchi AT, Haida M, Ohba H, Yamano M, Fukumoto D, et al. (2013) Liposome-encapsulated hemoglobin ameliorates ischemic stroke in nonhuman primates: Longitudinal observation. *Artif Organs* 37: 904-912.
- Kawaguchi AT, Haida M, Yamano M, Fukumoto D, Ogata Y, et al. (2010) Liposome-encapsulated hemoglobin ameliorates ischemic stroke in nonhuman primates: An acute study. *J Pharmacol Exp Ther* 332: 429-436.
- Kawaguchi AT, Kurita D, Furuya H, Yamano M, Ogata Y, et al. (2009) Liposome-encapsulated hemoglobin alleviates brain edema after permanent occlusion of the middle cerebral artery in rats. *Artif Organs* 33: 153-158.
- Komatsu H, Furuya T, Sato N, Ohta K, Matsuura A, et al. (2007) Effect of hemoglobin vesicle, a cellular-type artificial oxygen carrier, on middle cerebral artery occlusion- and arachidonic acid-induced stroke models in rats. *Neurosci Lett* 421: 121-125.
- Fukuta T, Ishii T, Asai T, Nakamura G, Takeuchi Y, et al. (2014) Real-time trafficking of PEGylated liposomes in the rodent focal brain ischemia analyzed by positron emission tomography. *Artif Organs* 38: 662-666.
- Ishii T, Asai T, Oyama D, Agato Y, Yasuda N, et al. (2013) Treatment of cerebral ischemia-reperfusion injury with PEGylated liposomes encapsulating FK506. *FASEB J* 27: 1362-1370.
- Ishii T, Asai T, Oyama D, Fukuta T, Yasuda N, et al. (2012) Amelioration of cerebral ischemia-reperfusion injury based on liposomal drug delivery system with asialo-erythropoietin. *J Control Release* 160: 81-87.
- Fukuta T, Ishii T, Asai T, Sato A, Kikuchi T, et al. (2015) Treatment of stroke with liposomal neuroprotective agents under cerebral ischemia conditions. *Eur J Pharm Biopharm* 97: 1-7.
- Imaizumi S, Woolworth V, Fishman RA, Chan PH (1990) Liposome-entrapped superoxide dismutase reduces cerebral infarction in cerebral ischemia in rats. *Stroke* 21: 1312-1317.
- Ishii T, Asai T, Fukuta T, Oyama D, Yasuda N, et al. (2012) A single injection of liposomal asialo-erythropoietin improves motor function deficit caused by cerebral ischemia/reperfusion. *Int J Pharm* 439: 269-274.

35. Joshi MD, Oesterling BM, Wu C, Gwizdz N, Pais G, et al. (2016) Evaluation of liposomal nanocarriers loaded with ETB receptor agonist, IRL-1620, using cell-based assays. *Neuroscience* 312: 141-152.
36. Fukuta T, Asai T, Sato A, Namba M, Yanagida Y, et al. (2016) Neuroprotection against cerebral ischemia/reperfusion injury by intravenous administration of liposomal fasudil. *Int J Pharm* 506: 129-137.
37. Kim H, Britton GL, Peng T, Holland CK, McPherson DD, et al. (2013) Nitric oxide-loaded echogenic liposomes for treatment of vasospasm following subarachnoid hemorrhage. *Int J Nanomedicine* 9: 155-165.
38. Peng T, Britton GL, Kim H, Cattano D, Aronowski J, et al. (2013) Therapeutic time window and dose dependence of xenon delivered via echogenic liposomes for neuroprotection in stroke. *CNS Neurosci Ther* 19: 773-784.
39. Luk YO, Chen WY, Wong WJ, Hu HH, Hsu LC, et al. (2004) Treatment of focal cerebral ischemia with liposomal nerve growth factor. *Drug Deliv* 11: 319-324.
40. Ramos-Cabrer P, Agulla J, Argibay B, Pérez-Mato M, Castillo J (2011) Serial MRI study of the enhanced therapeutic effects of liposome-encapsulated citicoline in cerebral ischemia. *Int J Pharm* 405: 228-233.
41. Suresh Reddy J, Venkateswarlu V, Koning GA (2006) Radioprotective effect of transferrin targeted citicoline liposomes. *J Drug Target* 14: 13-19.
42. Wang Z, Wu L, Xu M, Huang Y, Wang B, et al. (2015) Enhanced neuroprotection of ischemic stroke based on liposomal drug delivery system loading a novel uncoupler of ischemia-induced nNOS/PSD-95. *Current Signal Transd Ther* 10: 95-103.
43. Wang Z, Zhao Y, Jiang Y, Lv W, Wu L, et al. (2015) Enhanced anti-ischemic stroke of ZL006 by T7-conjugated PEGylated liposomes drug delivery system. *Scientific Reports*.
44. Yun X, Maximov VD, Yu J, Zhu H, Vertegel AA, et al. (2013) Nanoparticles for targeted delivery of antioxidant enzymes to the brain after cerebral ischemia and reperfusion injury. *J Cereb Blood Flow Metab* 33: 583-592.
45. Reddy MK, Labhasetwar V (2009) Nanoparticle-mediated delivery of superoxide dismutase to the brain: an effective strategy to reduce ischemia-reperfusion injury. *FASEB J* 23: 1384-1395.
46. Singhal A, Morris VB, Labhasetwar V, Ghorpade A (2013) Nanoparticle-mediated catalase delivery protects human neurons from oxidative stress. *Cell Death Dis* 4: e903.
47. Zhao G, Zang SY, Jiang ZH, Chen YY, Ji XH, et al. (2011) Postischemic administration of liposome-encapsulated luteolin prevents against ischemia-reperfusion injury in a rat middle cerebral artery occlusion model. *J Nutr Biochem* 22: 929-936.
48. Hwang H, Jeong HS, Oh PS, Na KS, Kwon J, et al. (2015) Improving cerebral blood flow through liposomal delivery of angiogenic peptides: Potential of 18F-fdg pet imaging in ischemic stroke treatment. *J Nucl Med* 56: 1106-1111.
49. Kim ID, Lim CM, Kim JB, Nam HY, Nam K, et al. (2010) Neuroprotection by biodegradable PAMAM ester (e-PAM-R)-mediated HMGB1 siRNA delivery in primary cortical cultures and in the postischemic brain. *J Control Release* 142: 422-430.
50. Omori N, Maruyama K, Jin G, Li F, Wang SJ, et al. (2003) Targeting of post-ischemic cerebral endothelium in rat by liposomes bearing polyethylene glycol-coupled transferrin. *Neurol Res* 25: 275-279.
51. Zhao H, Bao XJ, Wang RZ, Li GL, Gao J, et al. (2011) Postacute ischemia vascular endothelial growth factor transfer by transferrin-targeted liposomes attenuates ischemic brain injury after experimental stroke in rats. *Hum Gene Ther* 22: 207-215.
52. Lu YM, Huang JY, Wang H, Lou XF, Liao MH, et al. (2014) Targeted therapy of brain ischaemia using Fas ligand antibody conjugated PEG-lipid nanoparticles. *Biomaterials* 35: 530-537.
53. Bader KB, Bouchoux G, Peng T, Klegerman ME, McPherson DD, et al. (2015) Thrombolytic efficacy and enzymatic activity of rt-PA-loaded echogenic liposomes. *J Thromb Thrombolysis* 40: 144-155.
54. Britton GL, Kim H, Kee PH, Aronowski J, Holland CK, et al. (2010) *In vivo* therapeutic gas delivery for neuroprotection with echogenic liposomes. *Circulation* 122: 1578-1587.
55. Shaw GJ, Meunier JM, Huang SL, Lindsell CJ, McPherson DD, et al. (2009) Ultrasound-enhanced thrombolysis with tPA-loaded echogenic liposomes. *Thromb Res* 124: 306-310.
56. Smith DA, Vaidya SS, Kopeček JA, Huang SL, Klegerman ME, et al. (2010) Ultrasound-Triggered Release of Recombinant Tissue-Type Plasminogen Activator from Echogenic Liposomes. *Ultrasound Med Biol* 36: 145-157.
57. Petro M, Jaffer H, Yang J, Kabu S, Morris VB, et al. (2016) Tissue plasminogen activator followed by antioxidant-loaded nanoparticle delivery promotes activation/mobilization of progenitor cells in infarcted rat brain. *Biomaterials* 81: 169-180.
58. Tiebosch IA, Crielaard BJ, Bouts MJ, Zwartbol R, Salas-Perdomo A, et al. (2012) Combined treatment with recombinant tissue plasminogen activator and dexamethasone phosphate-containing liposomes improves neurological outcome and restricts lesion progression after embolic stroke in rats. *J Neurochem* 123: 65-74.
59. Agulla J, Brea D, Argibay B, Novo M, Campos F, et al. (2014) Quick adjustment of imaging tracer payload, for *in vivo* applications of the theranostic nanostructures in the brain. *Nanomedicine* 10: 851-858.
60. Agulla J, Brea D, Campos F, Sobrino T, Argibay B, et al. (2014) *In vivo* theranostics at the peri-infarct region in cerebral ischemia. *Theranostics* 4: 90-105.
61. Mouhieddine TH, Itani MM, Nokkari A, Ren C, Daoud G, et al. (2014) Nanotheragnostic Applications for Ischemic and Hemorrhagic Strokes: Improved Delivery for a Better Prognosis. *Curr Neurol Neurosci Rep* 15: 1-14.
62. Fisher M, Feuerstein G, Howells DW, Hurn PD, Kent TA, et al. (2009) Update of the Stroke Therapy Academic Industry Roundtable Preclinical Recommendations. *Stroke* 40: 2244-2250.